

KOPPREL 30 SL



DEFICIENCY CORRECTORS

Category: Deficiency Correctors

Formulation: Soluble Liquid (SL)



Foliar Application

KOPPREL 30 SL is formulated to maximise uptake and mobility of copper.

KOPPREL 30 SL is designed to penetrate the cuticle waxes of the plant.

KOPPREL 30 SL can also be applied via irrigation.

APPLICATION RATES

Crop	Concentration ml/ 100 L	Dose L/ha	Application
Grape	150 - 200	2.0	From budding to flower.
		1.0 - 1.5	Apply from bunch closure to harvest.
Mango, Avocado and Citrus fruits	150 - 200	2.0	Apply from the beginning of budding and during the fruit growth
Banana	200	1.5 - 2.0	Apply during fruit growth
Coffee	200	1.5 - 2.0	Apply during vegetative growth and fruit
Paprika, Peppers, and Tomato	150 - 200	1.0 - 1.5	Apply for 15 days after transplant until harvest.
Pumpkin, Melon, Watermelon	150 - 200	1.0 - 1.5	Apply 15 days after sowing or transplant to the neck of the plant.
Artichoke	150 - 200	1.0 - 1.5	Apply after transplantation and during vegetative growth
Potato	150 - 200	1.0 - 1.5	Apply during vegetative growth
Rice	150 - 200	1.0 - 1.5	Apply to tillering

GUARANTEED ANALYSIS

Element	% w/v
Copper (Cu)	7.6

pH: 3.40 / Density: 1.25

Packaging: 20 Litres

CONTACT US:

sales@biogrow.com.au

Melbourne: 03 9566 7252
W.A.: 0419 168 650
Shepparton: 0428 417 838

Mildura: 0419 250 192